

# *Agrobacterium*-mediated transformation of soybean and recovery of transgenic soybean plants

Paz, M., Martinez, J. C., Kalvig, A., Fonger, T., Wang, K. Improved cotyledonary node method using an alternative explant derived from mature seed for efficient *Agrobacterium*-mediated soybean transformation. *Plant Cell Reports*, 25: 206-213 (2006).

Paz, M., Wang, K. Soybean transformation and regeneration using half-seed explants. US Patent #7,473,822 (Issued January 6, 2009).

## Materials

### ⌘ Plasmid

Different plant transformation constructs that were derivatives of base vector pTF101.1 were introduced into *Agrobacterium tumefaciens* strain EHA101 (Hood et al. 1986). The base vector pTF101.1 is a derivative of the pPZP binary vector (Hajdukiewicz et al., 1994) that includes the right and left T-DNA border fragments from a nopaline strain of *A. tumefaciens*, a broad host origin of replication (pVS1) and a spectinomycin-resistant marker gene (*aadA*) for bacterial selection. The plant selectable marker gene cassette consists of (1) double 35S promoter (2x P35S) of cauliflower mosaic virus (CaMV) (Odell et al. 1985), (2) tobacco etch virus translational enhancer (Carrington and Freed 1990), (3) the phosphinothricin acetyl transferase (*bar*) gene from *Streptomyces hygroscopicus* that confers resistance to the herbicide phosphinothricin and its derivatives. pTF101.1 contains a multiple cloning site (MCS) for facilitating subcloning of the gene of interest in between the right border region and the plant selectable marker cassette. The soybean vegetative storage protein terminator (Mason et al., 1993) was cloned to the 3' end of the *bar* gene. The vector pTF102 was derived from pTF101.1 by inserting the P35S GUS intron cassette (Vancanneyt et al. 1990) into the *Hind* III site of pTF101.1. The *gus* gene contained a portable intron in its codon region (Vancanneyt et al., 1990) to prevent GUS activity in *Agrobacterium* cells.

### ⌘ Plant material

Soybean cultivars Thorne, Williams, and Williams 82

## Media

### ⌘ YEP Solid Medium

5 g/L Yeast extract, 10 g/L Peptone, 5 g/L NaCl<sub>2</sub>, 12 g/L Bacto-agar. pH to 7.0 with NaOH. Appropriate antibiotics should be added to the medium after autoclaving. Pour into sterile 100x15 plates (~25 ml per plate).

### ⌘ YEP Liquid Medium

5 g/L Yeast extract, 10 g/L Peptone, 5 g/L NaCl<sub>2</sub>. pH to 7.0 with NaOH. Appropriate antibiotics should be added to the medium prior to inoculation.

⌘ **Co-cultivation Medium**

1/10X Gamborg B5 Basal Medium (Contains: 1/10X B5 major salts, 1/10X B5 minor salts, 1/10 B5 vitamins, 2.8 mg/L Ferrous, 3.8 mg/L NaEDTA), 30 g/L Sucrose, 3.9 g/L MES, and 4.25 g/L Noble agar, pH 5.4. Filter sterilized GA<sub>3</sub> (0.25 mg/L), BAP (1.67 mg/L), Cysteine (400 mg/L), Dithiothrietol (154.2 mg/L), and 40 mg/L acetosyringone are added to this medium after autoclaving. Pour into sterile 100x15 mm plates (~88 plates/L). When solidified, overlay the co-cultivation medium with sterile filter paper to reduce bacterial overgrowth during co-cultivation (Whatman #1, 70 mm).

⌘ **Infection Medium**

1/10 X Gamborg B5 Basal Medium (Contains: 1/10X B5 major salts, 1/10X B5 minor salts, 1/10X B5 vitamins, 2.8 mg/L Ferrous, 3.8 mg/L NaEDTA), 30 g/L Sucrose, 3.9 g/L MES, pH 5.4. Filter sterilized GA<sub>3</sub> (0.25 mg/L), BAP (1.67 mg/L), and 40 mg/L acetosyringone are added to this medium after autoclaving.

⌘ **Shoot Induction Washing Medium (Optional)**

1X Gamborg B5 Basal Medium (Contains: 1X B5 major salts, 1X B5 minor salts, 1X B5 vitamins, 28 mg/L Ferrous, 38 mg/L NaEDTA), 30 g/L Sucrose, and 0.59 g/L MES, pH 5.7. Filter sterilized BAP (1.11 mg/L), Timentin (100 mg/L), Cefotaxime (100 mg/L), and Vancomycin (50 mg/L) are added to this medium after autoclaving.

⌘ **Shoot Induction Medium**

1X Gamborg B5 Basal Medium (Contains: 1X B5 major salts, 1X B5 minor salts, 1X B5 Vitamins, 28 mg/L Ferrous, 38 mg/L NaEDTA), 30 g/L Sucrose, 0.59 g/L MES, and 7 g/L Noble agar, pH 5.7. Filter sterilized BAP (1.11 mg/L), Timentin (50 mg/L), Cefotaxime (100 mg/L), Vancomycin (50 mg/L) and Glufosinate (8 mg/L) are added to this medium after autoclaving. Pour into sterile 100x20 mm plates (26 plates/L).

⌘ **Shoot Elongation Medium**

1X MS Modified Basal Medium with Gamborg Vitamins (Contains: 1X MS major salts, 1X MS minor salts, 1X B5 vitamins, 28 mg/L Ferrous, 38 mg/L NaEDTA, 30 g/L Sucrose, 0.59 g/L MES, and 7 g/L Noble agar, pH 5.7. Filter sterilized Asparagine (50 mg/L), L-Pyroglutamic Acid (100 mg/L), IAA (0.1 mg/L), GA<sub>3</sub> (0.5 mg/L), Zeatin-R (1 mg/L), Timentin (50 mg/L), Cefotaxime (100 mg/L), Vancomycin (50 mg/L), and Glufosinate (8 mg/L) are added to this medium after autoclaving. Pour into sterile 100x25 mm plates (22 plates/L).

⌘ **Rooting Medium**

½ X MS Modified Basal Medium with Gamborg Vitamins (Contains ½ X MS major salts, ½ X MS minor salts, ½ X B5 vitamins, 28 mg/L Ferrous, 38 mg/L NaEDTA), 20 g/L Sucrose, 0.59 g/L MES, and 7 g/L Noble agar, pH 5.6. Filter sterilized Indole-3-butyric acid (IBA, 1 mg/L), and glufosinate (3 mg/L) are added to this medium after autoclaving. Pour into sterile 150x25 mm vial (10 ml/vial).

## Methods

### ⌘ Seed Sterilization

1. Place mature soybean seeds in 100x15 mm Petri plates in a single layer (about 130 seeds per plate).
2. Arrange 3-4 plates in a bell jar desiccator within a fume hood in such a way that all interior plate surfaces are exposed and allow enough space to accommodate a 250 ml beaker.
3. Using appropriate hand protection, fill the 250 ml beaker with 100 ml of bleach and add 3.5 ml of concentrated (12 N) HCl drop wise along the side of the beaker.
4. Close the desiccator immediately and let stand overnight (16 hours).
5. After overnight exposure to chlorine gas, close the Petri plates and remove them to a laminar flow hood. Open the plates and allow them to air out for about 30 minutes to remove the excessive chlorine gas.

### ⌘ *Agrobacterium* Preparation

1. Bacteria cultures for weekly experiments are initiated from  $-80^{\circ}\text{C}$  glycerol stocks three days prior to an experiment. The vector system, pTF102 in EHA101, is cultured on YEP medium (An et al., 1988) containing 100 mg/L spectinomycin (for pTF102), 50 mg/L kanamycin (for EHA101), and 25 mg/L chloramphenicol (for EHA101).
2. 36 hours prior to the experiment start a 2 ml culture of *Agrobacterium* by inoculating a loop of bacteria from the fresh YEP plate in YEP liquid medium amended with antibiotics.
3. Allow the culture to grow to saturation (12-20 hours) at  $28^{\circ}\text{C}$  in a shaker incubator ( $\sim 250$  rpm). At the end of the day, transfer 0.25 ml of starter culture to a 1 L flask containing 250 ml of YEP medium amended with antibiotics.
4. Allow the culture to grow overnight at  $28^{\circ}\text{C}$ , 250 rpm to log phase ( $\text{OD}_{620} = 0.8-1.0$  for EHA101).
5. Collect *Agrobacterium* culture by pelleting at 4,000 rpm for 10 minutes at ambient temperature ( $\sim 22^{\circ}\text{C}$ ).
6. Resuspend the pellets in infection medium by pipetting through the pellet. Bacterial cell densities are adjusted to approximately  $\frac{1}{2}$  the original volume. A pellet from 50 ml of culture is resuspended to 25 mls with infection medium for approximate final density of  $0.5 \times 10^8$  cells/ml.
7. Gently shake the resulting infection medium at 60 rpm for at least 30 minutes before use.

### ⌘ Seed Imbibition

1. Under the laminar flow hood, approximately 20 hours prior to the infection experiment, add deionized sterile water to the sterilized seeds until the water is  $\frac{1}{4}$  cm from the top of the plate.
2. Completely cover plate with aluminum foil to block out light.

⌘ **Explant Preparation and Infection**

1. Remove aluminum foil from the imbibed soybean seeds. Transfer ~20 seeds to a sterile paper towel for dissection.
2. Using a #15 scalpel blade, make a longitudinal cut along the hilum to separate the cotyledons and remove the seed coat. Trim the embryonic axis found at the nodal end of the cotyledons to approximately 3 mm, and remove any remaining axial shoots/buds attached to the cotyledonary node.
3. Dissect 60 half-seed explants (30 seeds) into a 100 x 25 mm petri plate and add 30 ml of *Agrobacterium* infection media. Make sure the explants are completely covered by the infection media. Allow the explants to incubate at room temperature for 20-30 minutes with occasional gentle agitation.

⌘ **Co-Cultivation**

1. After infection, remove excess infection media by placing the seeds adaxial side up on sterile paper towel to drain excess liquid. Transfer half-seed explants to co-cultivation medium (9 per plate) so the flat, adaxial side is touching the filter paper.
2. Wrap the plates with parafilm or place plates in plastic bag with venting slits. Incubate the plates at 24°C under an 18:6 photoperiod (140  $\mu\text{moles s}^{-1} \text{m}^{-2}$ ) for 3 to 5 days.

⌘ **Shoot Induction**

1. After 3-5 days of co-cultivation, remove the half seeds from co-cultivation media and place briefly on sterile paper towel to remove excess liquid. Alternatively the half-seed explants can be washed briefly in shoot induction washing medium (~50 ml in a 100 x 25 sterile Petri plate, room temperature) then drained on sterile paper towel. Thirty half-seed explants may be washed for each plate of washing medium.
2. Place the explants on shoot induction medium (6 explants per plate). Half-seed explants should be oriented with the nodal end of the cotyledon imbedded in the medium and the regeneration region flush to the surface with flat side up at a 30°–45° angle. Transfer only half seeds containing intact trimmed embryonic axis.
3. Wrap each plate with vent tape and incubate at 24°C, 18:6 photoperiod for 14 days (if plates are stacked, by the end of the first week, rearrange each plate in each stack so that the top and bottom plates are switched and explants are exposed to light).
4. Explants should be transferred to fresh shoot induction medium after 14 days. Cut and discard shoots from the cot-node and apical area, and trim back the embryonic axis to 3 mm. Orient the tissue in such a way that the freshly cut surface is imbedded into the fresh shoot induction medium, with the differentiating region flush to the surface.
5. Maintain cultures in the Percival incubator under the same conditions described above for another 14 days.

⌘ **Shoot Elongation**

1. After 4 weeks (2 transfers) on shoot induction medium, remove the cotyledons from the explants and make a fresh cut at the base of the explant shoot pad flush to the medium.

2. Transfer the explants to fresh shoot elongation medium and incubate the tissue at 24°C, 18:6 photoperiod.
3. Transfer the tissue to fresh shoot elongation medium every 2 weeks. At each transfer make a fresh horizontal slice at the base of the shoot pad. Allow up to 8 weeks for shoot elongation.

#### ⌘ Rooting of Transgenic Plants

1. When shoots surviving glufosinate selection reach at least 3 cm, excise them from the shoot pad.
2. Transfer shoots to rooting medium in 150 x 25 mm glass vials with the stems of the shoots embedded approximately ½ cm into the media.
3. Incubate at 24°C, 18:6 photoperiod for 1-2 weeks.

#### ⌘ Plant Acclimatization and Liberty Screening

1. After 1-2 weeks, when the shoot develops more than two roots, transplant it into soil. Gently remove the plant from the rooting medium and wash off the roots with tap water to remove any excess medium.
2. Plants are placed in individual 2.5 inch plastic pot (cut from a four-pack) containing moistened Redi-Earth Peat-Lite soil mix (**Hummert Cat. # 10-2030-1**). Transplant plantlets to the soil and place them in a flat without holes, covered with a humidome. Allow the plantlets to grow at 24°C, 18:6 photoperiod for at least one week, watering as needed.
3. When the plantlets have at least two healthy trifoliates, an herbicide paint assay may be applied to confirm resistance to glufosinate. Using a transfer pipet, apply Liberty herbicide (250 mg/L), to the upper leaf surface along the midrib of two leaves on two different trifoliates. Score plantlets 5 days after painting. Resistant plantlets may be transplanted immediately to 1-gallon pots.
4. Fill 50% of a 1-gallon nursery pot with 4 drainage holes with Sunshine Universal Mix SB300. Add one tablet of Sierra 16-8-12 controlled release fertilizer with trace elements (7.5 g per 1-gal. pot) to each pot.
5. Add additional Universal Mix to bring the soil volume to 80% of the pot (~2" from top edge of pot). Too little does not hold enough water between watering, and too much does not allow enough water to be added at watering.
6. Transplant Liberty resistant plantlets, to the middle of the 1-gallon pot. The soil should cover all the roots. Be sure to plant the young plants deep enough or they will tip over when they grow taller.
7. Fill with water to the top edge of the pot. Let it drain completely, and water once more until the water reaches the top of the pot. After this time, water as needed.

#### ⌘ Plant Care

1. Marathon, for aphid control, may be added as part of the transplant step during summer months. If white flies or fungus gnats hover over pots, yellow sticky sheets may be used to reduce or eliminate the insects.

2. As plants grow, staking will be required to prevent plants from intertwining. Loosely bind elongated soybean branches to long bamboo stakes using twist ties.
3. Soybean pods on the same plant will dry at variable rates. To prevent pod shatter and consequent seed loss due to over drying, remove dry pods and store them in a paper bag until all pods on the plant are harvested.

## References

An G, Ebert PR, Mitra A, Ha SB (1988) Binary vectors. In: Gelvin SB and Schilperoort RA (eds) Plant Molecular Biology Manual. Kluwer Academic Publishers, Great Britain, pp 1-19.

Carrington JC, Freed DD (1990) Cap-independent enhancement of translation by a plant potyvirus 5' nontranslated region. J of Virology 64: 1590-1597.

Hajdukiewicz P, Svab Z, Maliga P (1994) The small, versatile pPZP family of *Agrobacterium* binary vectors for plant transformation. Plant Mol. Biol. 25:989-994.

Hood EE, Helmer GL, Fraley RT, Chilton M-D (1986) The hypervirulence of *Agrobacterium tumefaciens* A281 is encoded in a region of pTiBo542 outside of T-DNA. J. Bacteriol. 168:1291-1301.

Mason HS, DeWald D, Mullet JE (1993) Identification of a methyl jasmonate-responsive domain in the soybean vspB promoter. Plant Cell 5: 241-251.

Odell JT, Nagy F, Chua NH (1985) Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter. Nature 313, 810-812.

Olhoft PM, Somers DA (2001) L-cysteine increases *Agrobacterium*-mediated T-DNA delivery into soybean cotyledonary-node cells. Plant Cell Rep 20: 706-711.

Paz M, Martinez JC, Kalvig A, Fonger T, Wang K. (2006) Improved cotyledonary node method using an alternative explant derived from mature seed for efficient *Agrobacterium*-mediated soybean transformation. Plant Cell Reports, 25: 206-213.

Paz M, Wang K (2009) Soybean transformation and regeneration using half-seed explants. US Patent #7,473,822 (Issued January 6, 2009).

Vancanneyt G, Schmidt R, O'Connor-Sanchez A, Willmitzer L, Rocha-Sosa M (1990) Construction of an intron-containing marker gene: Splicing of the intron in transgenic plants and its use in monitoring early events in *Agrobacterium*-mediated plant transformation. Mol Gen Genet 220: 245-250.

Zhang Z, Aiqiu X, Staswick P, Clemente TE (1999) The use of glufosinate as a selective agent in *Agrobacterium*-mediated transformation of soybean. Plant Cell Tissue Org Cult 56: 37-46.